



LAB #: F000000-0000-0
 PATIENT: Sample Patient
 ID: P0000000000
 SEX: Female
 DOB: AGE: 55

CLIENT #: 12345
 DOCTOR:
 Doctor's Data, Inc.
 3755 Illinois Ave.
 St. Charles, IL 60174 U.S.A.

Comprehensive Parasitology, stool, x3

BACTERIOLOGY CULTURE

Expected/Beneficial flora

4+ Bacteroides fragilis group
 2+ Bifidobacterium spp.
 NG Escherichia coli
 2+ Lactobacillus spp.
 2+ Enterococcus spp.

 4+ Clostridium spp.
 NG = No Growth

Commensal (Imbalanced) flora

Dysbiotic flora

BACTERIA INFORMATION

Expected /Beneficial bacteria make up a significant portion of the total microflora in a healthy & balanced GI tract. These beneficial bacteria have many health-protecting effects in the GI tract including manufacturing vitamins, fermenting fibers, digesting proteins and carbohydrates, and propagating anti-tumor and anti-inflammatory factors.

Clostridia are prevalent flora in a healthy intestine. Clostridium spp. should be considered in the context of balance with other expected/beneficial flora. Absence of clostridia or over abundance relative to other expected/beneficial flora indicates bacterial imbalance. If *C. difficile* associated disease is suspected, a Comprehensive Clostridium culture or toxigenic *C. difficile* DNA test is recommended.

Commensal (Imbalanced) bacteria are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.

Dysbiotic bacteria consist of known pathogenic bacteria and those that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: consumption of contaminated water or food, exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

YEAST CULTURE

Normal flora

Dysbiotic flora

2+ Geotrichum spp

MICROSCOPIC YEAST

Result:

Mod

Expected:

None - Rare

Yeast in stool is expected at a level of none-rare. A microscopic finding of yeast in stool of few, moderate, or many may be helpful in identifying potential yeast overgrowth, or non-viable or dietary yeast.

YEAST INFORMATION

Yeast may normally be present in small quantities in the skin, mouth, and intestine. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast are not uniformly dispersed throughout the stool and this may lead to undetectable or low levels of yeast identified by microscopy, despite culture and identified yeast species. Conversely, microscopic examination may reveal a significant amount of yeast present but no viable yeast cultured. Yeast may not always survive transit through the intestines. Nonviable diet-derived yeast may also be detected microscopically. Consideration of clinical intervention for yeast detected microscopically should be made in the context of other findings and presentation of symptoms.

Comments:

Date Collected: 08/10/2017

Date Received: 08/15/2017

Date Reported: 08/24/2017

* *Aeromonas, Campylobacter, Plesiomonas, Salmonella, Shigella, Vibrio, Yersinia, & Edwardsiella tarda* have been specifically tested for and found absent unless reported.





LAB #: F000000-0000-0
PATIENT: Sample Patient
ID: P0000000000
SEX: Female
DOB: AGE: 55

CLIENT #: 12345
DOCTOR:
Doctor's Data, Inc.
3755 Illinois Ave.
St. Charles, IL 60174 U.S.A.

Comprehensive Parasitology, stool, x3

PARASITOLOGY/MICROSCOPY	PARASITOLOGY INFORMATION
<p>Sample 1 None Ova or Parasites Mod Yeast</p>	<p>Intestinal parasites are abnormal inhabitants of the gastrointestinal tract that have the potential to cause damage to their host. The presence of any parasite within the intestine generally confirms that the patient has acquired the organism through fecal-oral contamination. Damage to the host includes parasitic burden, migration, blockage and pressure. Immunologic inflammation, hypersensitivity reactions and cytotoxicity also play a large role in the morbidity of these diseases. The infective dose often relates to severity of the disease and repeat encounters can be additive.</p> <p>There are two main classes of intestinal parasites, they include protozoa and helminths. The protozoa typically have two stages; the trophozoite stage that is the metabolically active, invasive stage and the cyst stage, which is the vegetative inactive form resistant to unfavorable environmental conditions outside the human host. Helminths are large, multicellular organisms. Like protozoa, helminths can be either free-living or parasitic in nature. In their adult form, helminths cannot multiply in humans.</p> <p>In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and or blood, fever, nausea, or abdominal pain. However these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed or eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, and decreased immune function.</p> <p>In some instances, parasites may enter the circulation and travel to various organs causing severe organ diseases such as liver abscesses and cysticercosis. In addition, some larval migration can cause pneumonia and in rare cases hyper infection syndrome with large numbers of larvae being produced and found in every tissue of the body.</p> <p>One negative parasitology x1 specimen does not rule out the possibility of parasitic disease, parasitology x3 is recommended. This test is not designed to detect <i>Cyclospora cayetanensis</i> or <i>Microsporidia</i> spp.</p>
<p>Sample 2 None Ova or Parasites Rare Yeast</p>	
<p>Sample 3 None Ova or Parasites Rare Yeast</p>	

GIARDIA/CRYPTOSPORIDIUM IMMUNOASSAY			
	Within	Outside	Reference Range
<i>Giardia duodenalis</i>	Neg		Neg
<i>Cryptosporidium</i>	Neg		Neg

Giardia duodenalis (AKA *intestinalis* and *lamblia*) is a protozoan that infects the small intestine and is passed in stool and spread by the fecal-oral route. Waterborne transmission is the major source of giardiasis.

Cryptosporidium is a coccidian protozoa that can be spread from direct person-to-person contact or waterborne transmission.

Comments:

Date Collected: 08/10/2017
Date Received: 08/15/2017
Date Reported: 08/24/2017



LAB #: F000000-0000-0
 PATIENT: Sample Patient
 ID: P0000000000
 SEX: Female
 AGE: 55

CLIENT #: 12345
 DOCTOR:
 Doctor's Data, Inc.
 3755 Illinois Ave.
 St. Charles, IL 60174 U.S.A.


Yeast Susceptibilities: *Geotrichum spp*

NATURAL ANTIFUNGALS



Natural antifungal agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.

NON-ABSORBED ANTIFUNGALS



Non-absorbed antifungals may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative sensitivity is reported based upon the diameter of the zone of inhibition surrounding the disk.

AZOLE ANTIFUNGALS

	Resistant	S-DD	Susceptible
Fluconazole			S
Itraconazole			S
Ketoconazole			S

Susceptible results imply that an infection due to the fungus may be appropriately treated when the recommended dosage of the tested antifungal agent is used.
Susceptible - Dose Dependent (S-DD) results imply that an infection due to the fungus may be treated when the highest recommended dosage of the tested antifungal agent is used.
Resistant results imply that the fungus will not be inhibited by normal dosage levels of the tested antifungal agent.

Standardized test interpretive categories established for *Candida spp.* are used for all yeast isolates.

Comments:

Date Collected: 08/10/2017
 Date Received: 08/15/2017
 Date Completed: 08/24/2017

Yeast antifungal susceptibility testing is intended for research use only.
 Not for use in diagnostic procedures.

v10.11

INTRODUCTION

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific interpretive paragraphs are presented. If no significant abnormalities are found, interpretive paragraphs are not presented.

Clostridium spp

Clostridia are expected inhabitants of the human intestine. Although most clostridia in the intestine are not virulent, certain species have been associated with disease. Clostridium perfringens is a major cause of food poisoning and is also one cause of antibiotic-associated diarrhea. Clostridium difficile is a causative agent in antibiotic-associated diarrhea and pseudomembranous colitis. Other species reported to be prevalent in high amounts in patients with Autistic Spectrum Disorder include Clostridium histolyticum group, Clostridium cluster I, Clostridium bolteae, and Clostridium tetani.

If these disease associations are a concern further testing may be necessary.

Washington W, Allen S, Janda W, Koneman E, Procop G, Schreckenberger P, Woods, G. Koneman's Color Atlas and Textbook of Diagnostic Microbiology, 6th edition. Lippincott Williams and Wilkins; 2006. pg 931-939

Song Y, Liu C, Finegold SM. Real-Time PCR Quantitation of Clostridia in Feces of Autistic Children. Applied and Environmental Microbiology. Nov. 2004, 6459-6465.

Parracho H, Bingham MO, Gibson GR, McCartney AL. Differences Between the Gut Microflora of Children with Autistic Spectrum Disorders and That of Healthy Children. Journal of Medical Microbiology. 2005;54, 987-991.

Dysbiotic Yeast

Yeast was cultured from this stool specimen and the amount is considered to be dysbiotic. A positive yeast culture and sensitivity to prescriptive and natural agents is helpful in determining which anti-fungal agents to use as part of a therapeutic plan for chronic yeast syndrome. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast grows in colonies and is typically not uniformly dispersed throughout the stool. This may lead to undetectable or low levels of yeast identified by microscopy, despite a significant amount of yeast cultured.

Microscopic yeast

Microscopic examination has revealed yeast in this stool sample. The microscopic finding of yeast in the stool is helpful in identifying whether the proliferation of fungi, such as *Candida albicans*, is present. Yeast is normally found in very small amounts in a healthy intestinal tract. While small quantities of yeast (reported as none or rare) may be normal, yeast observed in higher amounts (few, moderate to many) is considered abnormal.

An overgrowth of intestinal yeast is prohibited by beneficial flora, intestinal immune defense (secretory IgA), and intestinal pH. Beneficial bacteria, such as *Lactobacillus* colonize in the intestines and create an environment unsuitable for yeast by producing acids, such as lactic acid, which lowers intestinal pH. Also, *Lactobacillus* is capable of releasing antagonistic substances such as hydrogen peroxide, lactocidin, lactobacillin, and acidolin.

Many factors can lead to an overgrowth of yeast including frequent use of antibiotics (leading to insufficient beneficial bacteria), synthetic corticosteroids, oral contraceptives, and diets high in sugar. Although there is a wide range of symptoms which can result from intestinal yeast overgrowth, some of the most common include brain fog, fatigue, recurring vaginal or bladder infections, sensitivity to smells (perfumes, chemicals, environment), mood swings/depression, sugar and carbohydrate cravings, gas/bloating, and constipation or loose stools.

A positive yeast culture (mycology) and sensitivity to prescriptive and natural agents is helpful in determining which anti-fungal agents to use as part of a therapeutic treatment plan for chronic colonic yeast. However, yeast are colonizers and do not appear to be dispersed uniformly throughout the stool. Yeast may therefore be observed microscopically, but not grow out on culture even when collected from the same bowel movement.

Beneficial Flora

One or more of the expected or beneficial bacteria are low in this specimen. Normally abundant include *Lactobacilli*, *Bifidobacteria*, *Clostridia*, *Bacteroides fragilis* group, enterococci, and some strains of *Escherichia coli*. The beneficial flora have many health-protecting effects in the gut, and as a consequence, are crucial to the health of the whole organism. Some of the roles of the beneficial flora include digestion of proteins and carbohydrates, manufacture of vitamins and essential fatty acids, increase in the number of immune system cells, break down of bacterial toxins and the conversion of flavinoids into anti-tumor and anti-inflammatory factors. *Lactobacilli*, *Bifidobacteria*, *Clostridia*, and enterococci secrete lactic acid as well as other acids including acetate, propionate, butyrate, and valerate. This secretion causes a subsequent decrease in intestinal pH, which is crucial in preventing an enteric proliferation of microbial pathogens, including bacteria and yeast. Many GI pathogens thrive in alkaline environments. *Lactobacilli* also secrete the antifungal and antimicrobial agents lactocidin, lactobacillin, acidolin, and hydrogen peroxide. The beneficial flora of the GI have thus been found useful in the inhibition of microbial pathogens, prevention and treatment of antibiotic associated diarrhea, prevention of traveler's diarrhea, enhancement of immune function, and

inhibition of the proliferation of yeast.

In a healthy balanced state of intestinal flora, the beneficial flora make up a significant proportion of the total microflora. Healthy levels of each of the beneficial bacteria are indicated by either a 3+ or 4+ (0 to 4 scale). However, some individuals have low levels of beneficial bacteria and an overgrowth of nonbeneficial (imbalances) or even pathogenic microorganisms (dysbiosis). Often attributed to the use of antibiotics, individuals with low beneficial bacteria may present with chronic symptoms such as irregular transit time, irritable bowel syndrome, bloating, gas, chronic fatigue, headaches, autoimmune diseases (e.g., rheumatoid arthritis), and sensitivities to a variety of foods. Treatment may include the use of probiotic supplements containing various strains of lactobacillus and bifidobacterium species and consumption of cultured or fermented foods including yogurt, kefir, miso, tempeh and tamari sauce. Polyphenols in green and ginseng tea have been found to increase the numbers of beneficial bacteria. If dysbiosis is present, treatment may also include the removal of pathogenic bacteria, yeast, or parasites.

Percival M. Intestinal Health. Clin Nutr In. 1997;5(5):1-6.

Fuller R. Probiotics in Human Medicine. Gut. 1991;32: 439-442.

Siitonen S, Vapaatalo H, Salminen S, et al. Effect of Lactobacilli GG Yoghurt in Prevention of Antibiotic Associated Diarrhea. Ann Med. 1990; 22:57-59.

Oksanen P, Salminen S, Saxelin M, et al. Prevention of Travelers' Diarrhea by Lactobacillus GG. Ann Med. 1990; 22:53-56.

Perdigon G, Alvarez M, et al. The Oral Administration of Lactic Acid Bacteria Increases the Mucosal Intestinal Immunity in Response to Enteropathogens. J Food Prot. 1990;53:404-410.

Valeur, N, et al. Colonization and Immunomodulation by Lactobacillus reuteri ATCC 55730 in the Human Gastrointestinal Tract. Appl Environ. Microbiol. 2004 Feb; 70(2):1176-81.

Elmer G, Surawicz C, and McFarland L. Biotherapeutic agents - a Neglected Modality for the Treatment and Prevention of Intestinal and Vaginal Infections. JAMA. 1996; 275(11):870-876.

Fitzsimmons N and Berry D. Inhibition of Candida albicans by Lactobacillus acidophilus: Evidence for Involvement of a Peroxidase System. Microbio. 1994; 80:125-133

Weisburger JH. Proc Soc Exp Biol Med 1999;220(4):271-5.